

CGMS-42 NASA-WP-01 22 May 2014

Prepared by NASA Agenda Item: D.2 Discussed in Plenary

NASA REPORT ON THE STATUS OF CURRENT AND FUTURE SATELLITE SYSTEMS

Executive summary

NASA currently operates 17 Earth Science missions. Although all missions were conceived as research missions, it has turned out that the efficiency of the communications and ground data handling systems has supported operational and near-real-time applications. All missions are currently producing data, but several also show signs of aging. Except for Suomi-NPP (Oct 2011), SAC-D/Aquarius (Jun 2011), LDCM/Landsat-8 (Feb 2013), and GPM Core (Feb 2014), all missions have passed their nominal design life, and are currently in extended operations.


NASA's Earth Science Program (\$1.8 Billion budget) is implementing a balanced and robust plan to accomplish a broad set of critical Earth observation measurements from space. The program advances knowledge of the integrated Earth system, the global atmosphere, oceans (including sea ice), land surfaces, ecosystems, and interactions between all elements, including the impacts of humans. A balance of satellite measurements, science research, technology development and applications are needed to address a complex global Earth system. NASA's plans include the launch of 13 missions and 7 instruments (on host missions) in the future.



NASA Report on the status of current and future satellite systems

1 CURRENT SATELLITE SYSTEMS


NASA currently operates 17 Earth Science missions (see Tables 1 and 2). All missions (except ACRIMSAT) are currently producing data, but several also show signs of aging. Except for Suomi-NPP (October 2011), SAC-D/Aguarius (June 2011), LDCM/Landsat-8 (February 2013), and GPM Core (February 2014), all missions have passed their nominal design life, and are currently in extended operations. Signs of battery aging have been observed in ACRIMSAT, QuikSCAT, TRMM, GRACE, CloudSat, and SORCE, all of which require intensive battery management and/or duty-cycling of instruments, which can reduce both quality and spatial/temporal coverage of the datasets. ACRIMSAT is currently unresponsive due to battery degradation, although recovery attempts continue. SORCE experienced a nearly 7-month data outage that ended in late February 2014 when it returned to limited operations during daylight only. (The Jason-1 mission failed in June 2013, and was decommissioned July 1, 2013). Instruments with reduced capability (noted in RED, Table 2) are the Landsat-7 ETM+ (failed Scan Line Corrector), QuikSCAT's SeaWinds (antenna no longer rotates so that the data are used primarily to crosscalibrate with other on-orbit scatterometers), Terra's ASTER (SWIR module is no longer functional), Aura's TES performs only Special Observations (no-longer performs Global Survey), Aqua's AMSU (Channels 4,5, and 7 are non-operational) and Aqua's AMSR-E (antenna rotates at 2rpm vs. 40rpm making the data useful for cross-calibration only). Instruments that no longer provide data (noted in RED, Table 2) are Aqua's HSB, TRMM's CERES, and Aura's HIRDLS. In addition, the TRMM VIRS, while functional, remains off to preserve battery life. All other sensors are fully functional and are producing standard products that meet or exceed specifications.



The most recently launched NASA mission is the The Global Precipitation Measurement (GPM) Core Observatory, launched on February 27, 2014. The GPM mission will help advance our understanding of Earth's water and energy cycles, improve the forecasting of extreme events that cause natural disasters, and extend current capabilities of using satellite precipitation information to directly benefit society. The GPM mission will provide unprecedented and freely available data on rain and snowfall from its GPM Microwave Imager (GMI) and Dual-frequency Precipitation Radar (DPR) instruments. The GMI image on the left is an extra-tropical cyclone off the coast of Japan on March 10, 2014. Red areas indicate heavy



precipitation, while yellow and blue areas indicate less intense precipitation. Once new missions are launched, NASA must continually monitor their positions to avoid collisions with other satellites. Increased solar activity has led to more uncertainty in collision analysis calculations and consequently intensified analysis and planning activities to determine collision avoidance maneuvers. A history of collision avoidance maneuvers is shown in the figure below. Active monitoring of close approach events has steadily increased since 2008. In addition, potential conjunctions between operational, maneuverable satellites have increased, necessitating communication between the satellite operators in order to coordinate avoidance maneuver planning. In addition to increasing the resources dedicated to collision assessment, NASA continually improves the agency's orbital debris procedures, and invests in analysis tool improvements.



1.1 Research Missions for Operational Use

Although all missions were conceived as research missions, it has turned out that the efficiency of the communications and ground data handling systems has supported operational and near-real-time applications. Our interagency partners have rated most NASA missions as High Utility for operational applications, with Terra, Aqua, TRMM and Suomi-NPP rated Very High. All missions have met their original success criteria and are meeting the objective for sustained measurements on decadal time scales. This objective is met not only due to the satellites' longevity, but also to the sustained calibration/validation program and the data systems tools which enhance data quality and access. Continued operation of the missions is determined through a biennial



science review process, called the "Senior Review", which evaluates the continuing science value. Operational uses of the missions are considered in the review, but science remains the defining factor for continuation. The most recent Senior Review, conducted March-May 2013, found that all of NASA's missions currently in extended operations are still producing valuable science datasets for research, and should be extended for another 2 years. The next Senior Review is scheduled for mid-2015.

1.2 Real-time Direct Broadcast Data

Several of NASA's missions provide for the real-time transmission of satellite data to the ground in support of operational activities and disaster monitoring. As the Earth is being observed by satellite instruments on these platforms the data is transmitted using omnidirectional antennas. Users who have compatible ground receiving equipment and are in direct line of sight to the satellite may receive these transmissions. This Direct Broadcast capability is currently available for selected instruments on the Aqua, Terra, and Suomi-NPP missions. More information on the required hardware and ground station processing software can be found at NASA's Direct Readout Laboratory (DRL) website: http://directreadout.sci.gsfc.nasa.gov. NASA also provides access to Near Real-Time (NRT) global data and products from the MODIS (on Terra and Aqua), OMI and MLS (on Aura), and AIRS (on Aqua) instruments in less than 2.5 hours from observation from the Land and Atmosphere Near real-time Capability for EOS (LANCE) data system at http://earthdata.nasa.gov/lance.



Table 1 - Current NASA LEO Satellites

| Satellite | Operator | Equatorial Crossing Time | Mean Altitude | Launch Date | Data Access | Instrument Status |
|--|-----------------------------|--------------------------------|------------------|-----------------|--|--|
| Jason-2 (Op) (Ocean Surface Topography Mission) | NASA/NOAA, EUMETSAT/CNES | 66-deg Non Sun-Sync | 1336 | 20-Jun- 2008 | <u>Handbook</u> | Science: Sea surface topography (Follow on to Jason-1) Instruments: LRA, DORIS, POSEIDON-3, AMR, GPSP |
| Suomi-NPP (Op) | USA NASA/NOAA | 13:30 | 833 | 28-Oct- 2011 | <u>Suomi Data</u> <u>Direct</u> <u>Broadcast</u> | Science: Atmospheric dynamics, water and energy cycle, clouds and aerosols, radiation, GHG, air/sea fluxes; also supporting operational weather forecasting & ozone monitoring Instruments: CrIS, CERES, VIIRS, ATMS, OMPS |



Table 2 - Current NASA Research and Development (R&D) Satellites

| Satellite | Space Agency | Equatorial Crossing Time and Mean Altitude | Launch Date | Data Access | Instruments | Status, Applications and Other Information |
|-----------|-----------------|---|----------------|---------------------------------------|--|---|
| TRMM | NASA/JAXA | 35 Deg Inclination Non Sun-Sync 402 km | 28-Nov-1997 | <u>PMM Data</u> <u>TRMM Data</u> | LIS, PR, <mark>CERES, VIRS</mark> , TMI | Atmospheric dynamics, water and energy cycle, lightning, precipitation, radiation |
| Landsat-7 | NASA/USGS | 10:05 (D) 705 km | 15-Apr-1999 | Earth Explorer | ETM+ | Earth resources, land surface, environmental and disaster monitoring, agriculture and forestry, ice and snow cover |
| QuikSCAT | NASA | 6:00 (A) 803 km | 19-Jun-1999 | PO.DAAC | SeaWinds | Sea surface wind vectors |
| Terra | NASA | 10:30 (D) 705 km | 18-Dec-1999 | <u>Terra Data</u> Direct Broadcast | ASTER, MODIS, MOPITT, MISR, CERES | Atmospheric dynamics and chemistry, water and energy cycle, clouds, aerosols, radiation, GHG, carbon and water, air-land exchange |
| ACRIMSAT | NASA | 10:50 (D) 720 km | 20-Dec-1999 | <u>ASDC</u> | ACRIM-III | Total solar irradiance, solar constant |



| NMP EO-1 | NASA | 9:45 (D) 680 km | 21-Nov-2000 | Archive Earth Explorer New Data | ALI, Hyperion, LEISA AC | Land surface and earth resources |
|--------------------|-----------|---|-------------|---------------------------------------|--|---|
| GRACE | NASA/DLR | 89 Deg Inclination Non Sun-Sync 485 km | 17-Mar-2002 | PO.DAAC | MWA, Accelerometers, GPS | Earth mass distribution, with application to ground water, ocean currents and ice sheets, GPS (P,T,humidity) |
| Aqua (EOS PM-1) | NASA | 13:30 (A) 705 km | 4-May-2002 | EOSDIS Direct Broadcast | MODIS, AIRS, CERES, AMSU-A, AMSR-E, HSB | Atmospheric dynamics, water and energy cycle, clouds and aerosols, radiation, GHG, air/sea fluxes, precipitation |
| SORCE | NASA | 40 Deg Inclination Non Sun-Sync 640 km | 25-Jan-2003 | DISC | SIM, SOLSTICE, TIM, XPS | Total and spectral solar irradiance |
| Aura | NASA | 13:45 (A) 705 km | 15-Jul-2004 | DISC | MLS, TES, <mark>HIRDLS,</mark> OMI | Chemistry and dynamics of atmosphere, O3, GHG, aerosols |
| CALIPSO | NASA/CNES | 13:30 (A) 705 km | 28-Apr-2006 | <u>ASDC</u> | CALIOP, IIR, WFC | Aerosols and clouds |
| CloudSat | NASA/CSA | 13:30 (A) 705 km | 28-Apr-2006 | Cloudsat DPC | CPR | Cloud vertical profiling |



| SAC-D / Aquarius | NASA/CONAE | 18:00 (A) 651 km | 10-Jun-2011 | PO.DAAC | L-Band Radiometer, L-Band Scatterometer, CARMEN-1, DCS, HSC, Lagrange, MWR, NIRST, ROSA, SODAD, TDP | Sea Surface Salinity |
|---------------------|------------|---|-------------|----------------------------|--|---|
| LDCM (Landsat-8) | NASA/USGS | 10:05 (D) 705 km | 11-Feb-2013 | Landsat-8 Data Products | OLI, TIRS | Earth resources, land surface, environmental and disaster monitoring, agriculture and forestry, ice and snow cover |
| GPM Core | NASA/JAXA | 65 Deg Inclination Non Sun-Sync 407 km | 27-Feb-2014 | PMM Data | GMI, DPR | Global precipitation, evaporation, water cycle |

Failed Instruments

* CERES on TRMM, HSB on Aqua, and HIRDLS on Aura

Reduced Function Instruments

- * SeaWinds on QuikSCAT (no antenna rotation, only used for cross-calibration)
- * ETM+ on Landsat-7 (failed scan line corrector)
- * ASTER (SWIR module not functioning)
- * AMSU on Aqua (channel-4 failed)
- * AMSR-E on Aqua (reduced rotation rate for cross-calibration with AMSR-2)
- * SORCE Battery degradation, Instruments turned off during orbit night, data retrieved for up to 4 orbits/day only
- * Cloudsat Battery degradation, Instruments turned off during orbit night
- * VIRS on TRMM (functional but turned off to preserve battery)



2 FUTURE SATELLITE SYSTEMS

With the U.S. President's FY2015 budget request (\$1.77 Billion), NASA's Earth Science Program is implementing a balanced and robust plan to accomplish a broad set of critical Earth observation measurements from space. The program advances knowledge of the integrated Earth system, the global atmosphere, oceans (including sea ice), land surfaces, ecosystems, and interactions between all elements, including the impacts of humans. A balance of satellite measurements, science research, technology development and applications are needed to address a complex global Earth system. Table 3 summarize NASA's future plans for the launch of 13 missions and 7 instruments (on host missions).

2.1 Earth Systematic Missions (ESM)

NASA's ESM includes a broad range of multi-disciplinary science investigations aimed at developing a scientific understanding of the Earth system and its response to natural and human-induced forces and changes. The ESM program develops Earth observing research satellite missions, manages the operation of NASA facility research missions once on orbit, and produces standard mission products in support of NASA and National research, applications, and policy communities. The five current flight missions in formulation or development contained in the ESM program are the Ice, Cloud, and Land Elevation Satellite (ICESat)-2, Soil Moisture Active-Passive (SMAP), Stratospheric Aerosol and Gas Experiment (SAGE)-III, Gravity Recovery and Climate Experiment Follow-On (GRACE-FO) and Surface Water Ocean Topography (SWOT) missions.

The program has adjusted the mission timelines and budgets to accommodate increasing launch vehicle costs (for SMAP) and the reallocation of funding based on Agency priorities. The SMAP launch date and funding profile is driven by launch vehicle availability and the current plan supports an October 2014 launch. The Agency continues with the pre-formulation studies, formulation, and development of other Decadal and climate missions such as NASA ISRO-Synthetic Aperture Radar (NI-SAR), Pre-Aerosols, Carbon and Ecosystems (PACE), Climate Absolute Radiance and Refractivity Observatory (CLARREO), Active Sensing of Carbon dioxide Emissions over Nights, Days and Seasons (ASCENDS), Aerosols, Clouds and Ecosystems (ACE), Geostationary Coastal and Air Pollution Events (GEO-CAPE), and Hyperspectral Infrared Imager (HyspIRI).

2.2 Earth System Science Pathfinder (ESSP)

ESSP provides an innovative approach to Earth science research by providing frequent, regular, competitively selected opportunities that accommodate new and emerging scientific priorities and measurement capabilities. These opportunities represent a series of relatively low-to-moderate cost, small-to-medium sized missions. They are competitively selected, principal investigator led missions that



focus on scientific objectives to support a selected subset of studies of the atmosphere, oceans, land surface, polar ice regions, or solid Earth. NASA currently funds the development of the Orbiting Carbon Observatory (OCO-2) mission, the Earth Venture-class missions, and several other missions and instruments in pre-formulation under ESSP, including the OCO-3 instrument. The current projected OCO-2 launch date is July 2014.

The Earth Venture-class (EV) missions are part of a competitive program to select small instruments, small satellites, or airborne science campaigns to complement the strategic NASA Earth science missions. In FY 2014, NASA will continue the five airborne science investigations selected through the initial Venture Class solicitation (EVS-1) in FY 2009 and started in FY 2010. NASA selected the Cyclone Global Navigation Satellite System (CYGNSS) mission as part of the Earth Venture Mission (EVM-1) solicitation in 2012. CYGNSS is currently in development and will launch in 2016. The Tropospheric Emissions: Monitoring of Pollution (TEMPO) instrument was selected in November 2012 as part of the Earth Venture Instrument (EVI-1) solicitation. TEMPO will be mounted on a commercial communications satellite in geostationary orbit to monitor air pollutants over North America no earlier than 2018. Future solicitations for the Earth Venture Class projects will be released every 4 years for the EVS and EVM elements and approximately every 18-months for EVI. The next planned release is EVM-2 and EVI-3 in 2015.



Table 3 - Future NASA Research and Development (R&D) Satellites

| Satellite | Space Agency | Equatorial Crossing Time and Mean Altitude | Launch Date | Data Access | Instruments | Status, Applications and Other Information |
|---|-----------------|---|-------------|----------------|------------------------------------|--|
| OCO-2 | NASA | 13:30 (A) 705 km | July 2014 | | Spectrometer | Carbon Dioxide sources and sinks |
| SAGE-III-ISS Intl. Space Station Instrument only | NASA | 51.6 Deg Inclination Non Sun-Sync 407 km | TBD | | Spectrometer | Stratospheric ozone, aerosols, and water vapor |
| CATS-ISS Intl. Space Station Instrument only | NASA | 51.6 Deg Inclination Non Sun-Sync 407 km | Aug 2014 | | LIDAR | Atmospheric pollution, dust, smoke, and aerosols |
| RapidScat-ISS Intl. Space Station Instrument only | NASA | 51.6 Deg Inclination Non Sun-Sync 407 km | Aug 2014 | | Scatterometer | Ocean surface wind speed and direction |
| SMAP | NASA | 18:00 (A) 685 km | Nov 2014 | | L-Band Radar, L-Band Radiometer | Soil Moisture, Freeze-thaw state |
| LIS-ISS Intl. Space Station Instrument only | NASA | 51.6 Deg Inclination Non Sun-Sync 407 km | Feb 2016 | | Optical Imager | Lightning |



| CYGNSS | NASA | 8 small satellites, 35 Deg Inclination, Non Sun-Sync 500 km | Oct 2016 | | GPS | Ocean surface winds for tropical storms and hurricanes. |
|---|-----------|--|---------------------|---------|---|--|
| ICESat-II | NASA | 92 Deg Inclination Non Sun-Sync 478 km | Oct 2017 | | ATLAS | Ice sheet thickness, sea ice thickness, vegetation height, carbon and biomass |
| OCO-3-ISS Intl. Space Station Instrument only | NASA | 51.6 Deg Inclination Non Sun-Sync 407 km | TBD | | Spectrometer | Carbon Dioxode sources and sinks |
| HICO-ISS Intl. Space Station Instrument only | NASA | 51.6 Deg Inclination Non Sun-Sync 407 km | TBD | | Imaging Spectrometer | Coastal ocean water clarity, bottom types, bathymetry, and on-shore vegetation |
| GRACE-FO (Follow-On) | NASA/GFZ | 89 Deg Inclination Non Sun-Sync 490 km | Aug 2017 | | Gravity, GPS | Ocean currents and mass, ice sheets, GPS (Pressure, Temperature, Humidity) |
| TEMPO Hosted Payload Instrument only | NASA | Geosynchronous | <u>>Nov</u> 2018 | | UV and VIS Spectrometer | Atmospheric pollution over North America. Tropospheric ozone, ozone precursors, aerosols, and clouds. |
| SWOT | NASA/CNES | 78 Deg Inclination Non Sun-Sync 891 km | 2020 | PO.DAAC | Ka-Band Radar Interferometer, AMR, GPSP, LRA, Poseidon Altimeter | Oceanography (wide swath ocean surface topography) and Hydrology (lake levels, river discharge) |
| PACE (Pre-ACE) | NASA | Sun-Synchronous 650 km | <u>></u> 2020 | | Spectrometer, Polarimeter | Aerosols, ocean color |



| ASCENDS | NASA | 10:30 (A) 450 km | <u>></u> 2023 | Laser | Carbon Dioxide (day and night) |
|----------|------|---|------------------|---|--|
| CLARREO | NASA | 90 Deg Inclined | TBD | IR, RS, GNSS | Spectrally resolved and calibrated Infrared (IR) and Reflected Solar (RS) Earth radiance, GNSS (T,P,humidity) |
| NI-SAR | NASA | 98 Deg Inclination, Sun-Sync (6AM- 6PM), 747 km | 2020 | L-band, S-band SAR (repeat-pass interferometry, polarimetry) | Earth surface deformation, ecosystems and biomass change, ice motion |
| HyspIRI | NASA | TBD | TBD | Hyperspectral and TIR Imagers | Terrestrial and aquatic ecosystems, fires, mineral resources, volcanoes |
| GEO-CAPE | NASA | Geosynchronous | TBD | UV-Vis-NIR, IR imagers (CO detection) | Air pollution forecasting and transport, sources of aerosols and O3, coastal ecosystems, CO, NO2, SO2, HCHO |
| ACE | NASA | TBD | TBD | Spectrometer, Polarimeter, LIDAR, Cloud Radar | Aerosols, ocean color, cloud profiles |